



UTA Patient Measles, Mumps, and Rubella Vaccination Form

SCREENING FOR ADULT IMMUNIZATION TO BE COMPLETED BY PATIENT

Patient Name: _____ UTA ID#: _____

DOB: _____ Phone #: _____

Please answer the questions with a yes or no. If the question is not clear, please ask a nurse to explain it.

- 1) Are you breastfeeding, pregnant or is there a chance you are going to become pregnant in the next **2 months**?
- 2) Are you sick today?
- 3) Have you ever had a serious reaction after receiving a vaccination?
- 4) History of Gullain-Barre Syndrome within 6 weeks after receiving a flu vaccine?
- 5) Do you have cancer, Guillan-Barre Syndrome, AIDS or any other immune system problems?
- 6) Are you taking any steroids or anticancer drugs, OR are you currently undergoing radiation?
- 7) During the past year have you received a transfusion of blood or plasma or been given medicine called Immune globulin?
- 8) Do you have a condition that makes you bleed or bruise easily?
- 9) Do you have a severe allergy to gelatin or neomycin?
- 10) **Have you received any vaccine in the past 4 weeks?**

Statement of Authorization

I authorize UTA Health Center personnel or designee to administer the immunization and any necessary subsequent treatment. I understand I am ultimately responsible for any charges incurred during my treatment. I received a copy of the Vaccine Information Sheet (VIS) for the vaccine. I understand the benefits and risks of the vaccine, and that the CDC recommends you wait at the place you receive your injection for 20 minutes after injection.

UTA ID # of Person Receiving Vaccine

Date

MMR Vaccine (Measles, Mumps, and Rubella): *What You Need to Know*

Many Vaccine Information Statements are available in Spanish and other languages. See www.immunize.org/vis

Hojas de información sobre vacunas están disponibles en español y en muchos otros idiomas. Visite www.immunize.org/vis

1 Why get vaccinated?

MMR vaccine can prevent **measles, mumps, and rubella**.

- **MEASLES (M)** can cause fever, cough, runny nose, and red, watery eyes, commonly followed by a rash that covers the whole body. It can lead to seizures (often associated with fever), ear infections, diarrhea, and pneumonia. Rarely, measles can cause brain damage or death.
- **MUMPS (M)** can cause fever, headache, muscle aches, tiredness, loss of appetite, and swollen and tender salivary glands under the ears. It can lead to deafness, swelling of the brain and/or spinal cord covering, painful swelling of the testicles or ovaries, and, very rarely, death.
- **RUBELLA (R)** can cause fever, sore throat, rash, headache, and eye irritation. It can cause arthritis in up to half of teenage and adult women. If a woman gets rubella while she is pregnant, she could have a miscarriage or her baby could be born with serious birth defects.

Most people who are vaccinated with MMR will be protected for life. Vaccines and high rates of vaccination have made these diseases much less common in the United States.

2 MMR vaccine

Children need 2 doses of MMR vaccine, usually:

- First dose at 12 through 15 months of age
- Second dose at 4 through 6 years of age

Infants who will be traveling outside the United States when they are between 6 and 11 months of age should get a dose of MMR vaccine before travel. The child should still get 2 doses at the recommended ages for long-lasting protection.

Older children, adolescents, and adults also need 1 or 2 doses of MMR vaccine if they are not already immune to measles, mumps, and rubella. Your

health care provider can help you determine how many doses you need.

A third dose of MMR might be recommended in certain mumps outbreak situations.

MMR vaccine may be given at the same time as other vaccines. Children 12 months through 12 years of age might receive MMR vaccine together with varicella vaccine in a single shot, known as MMRV. Your health care provider can give you more information.

3 Talk with your health care provider

Tell your vaccine provider if the person getting the vaccine:

- Has had an **allergic reaction after a previous dose of MMR or MMRV vaccine**, or has any **severe, life-threatening allergies**.
- Is **pregnant**, or thinks she might be pregnant.
- Has a **weakened immune system**, or has a **parent, brother, or sister with a history of hereditary or congenital immune system problems**.
- Has ever had a **condition that makes him or her bruise or bleed easily**.
- Has recently had a **blood transfusion or received other blood products**.
- Has **tuberculosis**.
- Has **gotten any other vaccines in the past 4 weeks**.

In some cases, your health care provider may decide to postpone MMR vaccination to a future visit.

People with minor illnesses, such as a cold, may be vaccinated. People who are moderately or severely ill should usually wait until they recover before getting MMR vaccine.

Your health care provider can give you more information.



4 Risks of a vaccine reaction

- Soreness, redness, or rash where the shot is given and rash all over the body can happen after MMR vaccine.
- Fever or swelling of the glands in the cheeks or neck sometimes occur after MMR vaccine.
- More serious reactions happen rarely. These can include seizures (often associated with fever), temporary pain and stiffness in the joints (mostly in teenage or adult women), pneumonia, swelling of the brain and/or spinal cord covering, or temporary low platelet count which can cause unusual bleeding or bruising.
- In people with serious immune system problems, this vaccine may cause an infection which may be life-threatening. People with serious immune system problems should not get MMR vaccine.

People sometimes faint after medical procedures, including vaccination. Tell your provider if you feel dizzy or have vision changes or ringing in the ears.

As with any medicine, there is a very remote chance of a vaccine causing a severe allergic reaction, other serious injury, or death.

5 What if there is a serious problem?

An allergic reaction could occur after the vaccinated person leaves the clinic. If you see signs of a severe allergic reaction (hives, swelling of the face and throat, difficulty breathing, a fast heartbeat, dizziness, or weakness), call **9-1-1** and get the person to the nearest hospital.

For other signs that concern you, call your health care provider.

Adverse reactions should be reported to the Vaccine Adverse Event Reporting System (VAERS). Your health care provider will usually file this report, or you can do it yourself. Visit the VAERS website at www.vaers.hhs.gov or call **1-800-822-7967**. *VAERS is only for reporting reactions, and VAERS staff do not give medical advice.*

6 The National Vaccine Injury Compensation Program

The National Vaccine Injury Compensation Program (VICP) is a federal program that was created to compensate people who may have been injured by certain vaccines. Visit the VICP website at www.hrsa.gov/vaccinecompensation or call **1-800-338-2382** to learn about the program and about filing a claim. There is a time limit to file a claim for compensation.

7 How can I learn more?

- Ask your healthcare provider.
- Call your local or state health department.
- Contact the Centers for Disease Control and Prevention (CDC):
 - Call **1-800-232-4636 (1-800-CDC-INFO)** or
 - Visit CDC's www.cdc.gov/vaccines

Vaccine Information Statement (Interim)
MMR Vaccine



Office use only